

INTERNATIONAL JOURNAL OF ADVANCED RESEARCH IN BIOLOGICAL SCIENCES

(ISSN:2348-8069)



[Home](#) | [Editorial Board](#) | [Guidelines to Authors](#) | [Submission of Papers](#) |
[Indexing and Abstracting](#) |

[Current Issue](#) | [Archive](#) | [Join Reviewer and Editorial Member](#) | [Contact Us](#)

IJARBS

International Journal of Advanced Research in Biological Sciences publishes original papers, review articles, and short communications. The mission of International Journal of Current Research in Biological Sciences is to significantly broaden the knowledge base of its readers and in this sense; the journal shall focus on only those papers that fall within its scope. The scope of the journal encompasses research articles, original research reports, reviews, short communications and scientific commentaries in the fields of science and Agricultural sciences and medical science and other fields of basic and applied sciences.

International Journal of Advanced Research in Biological Sciences aims to publishers various fields of biological sciences including Life sciences, Medical sciences, Microbiology, Biotechnology, Agricultural Sciences, Chemical Sciences, and Environmental Sciences, Botany, Plant Science, Biochemistry, Zoology, Marine Biology, Genetics, Molecular Biology, Pharmacogenomics, Pharmacology, Medicine and Dentistry, Veterinary Science, Anatomy, Nutrition and Metabolism, Cancer Research and Experimental Oncology, Clinical Pathology and Forensic, Medicine, Agricultural Engineering, Agricultural Biotechnology, Agricultural Microbiology and Plant Pathology.



VOL. 2. ISSUE – 6 (June 2015)

S.No	Articles	Page No	Download
1.	Effects of Silicon on membrane characteristics, photosynthetic pigments, antioxidative ability, and mineral element contents of faba bean (<i>Vicia faba</i> L.) plants grown under Cd and Pb stress. Sharifa S. Abu-Muriefah	01-17	
2.	Retinol Binding Protein-4: a Novel Predictor of Insulin Resistance in Lean Subjects Hosny SS, El-Sherbeny AA, Elewa AA, Kamal El -dine MM	18-26	

3.	Antimicrobial activity of sea anemone <i>Stichyodactyla hadonii</i> and <i>Anthopleura elegantissima</i> extracts against human pathogens Shani.T.John, S. Velmurugan, D. Senthil Nagaraj S.kumaran, S.R.Pugazhvendan	27–35	
4.	Identification of MDR <i>M.tuberculosis</i> and corresponding changes in <i>rpoB</i> gene S.Asokan and J.Sangeetha	36–44	
5.	Probiotics in Aquaculture Madhura Mukadam	45–47	
6.	GeneXpert System for Detection of Staphylococci from Blood Cultures and their Resistance to Methicillin (our first experience) Zaklina Cekovska, Kaftandzieva A., Petrovska M., Trajkovska Dokic E., Jankoska G., Kotevska V., Ristovska N., Panovski N.	48–54	
7.	Feeding selection by Timor deer (<i>Cervus timorensis</i> Blainville) in relation to chemical and physical properties of the forage plants at Bali Barat National Park I Ketut Ginantra, I Wayan Suarna, I Wayan Kasa and Deden Ismail	55–65	
8.	Amylase production from <i>Aspergillus flavus</i> using millet pomace by Solid State Fermentation (SSF) Famotemi, A. C., Lawal, A. K., Esan A. O., Salihu, I., Tijani, A. A. and Dike E. N.	66–72	
9.	Abd El-hadi EA El-sherbeny AA, Baha El-din AM , Shaaban MA , Badran HA, Balata MG	73–81	
10.	Dr. Batool Hassan Al-Ghurabi; Dr. Ahmed Abdulhassan Abbas; Dr. Mustafa M. Salah; Dr. Fatin M. Jarallah	82–85	
11.	Paramita Biswas and Rati Kanta Ghosh	86–92	
12.	Eskandari E., Shayestehfar A. and Noori M.	93–97	
13.	BhavyaKavithaDwarapureddi and VaraSaritha	98–113	
14.	Hany Haroun Kaisar, MD; Gamal Fawzy, MD; Ahmed A Darwish, MD	114–120	
15.	<i>E.coli</i> strains isolated from clinical samples of Cuddalore district, Tamilnadu. R.S.Vigneshwari and G. Thiruneelakandan	121–128	

Research Article



Feeding selection by Timor deer (*Cervus timorensis* Blainville) in relation to chemical and physical properties of the forage plants at Bali Barat National Park

***I Ketut Ginantra⁽¹⁾, I Wayan Suarna⁽²⁾, I Wayan Kasa⁽¹⁾ and Deden Ismail⁽³⁾**

⁽¹⁾Department of Biology, Faculty of Mathematic and Natural Science and ⁽²⁾Department of Nutrition and Forage Science, Faculty of Husbandry, Udayana University, Bali-Indonesia

⁽³⁾Department of Biology, Faculty of Mathematic and Natural Science, Mahasaraswati University, Bali-Indonesia

*Corresponding author

Abstract

This study was conducted to determine the relationship feeding selection of timor deer (*Cervus timorensis*) by a factor of the chemical and physical properties of forage available in the habitat. Composition of plants in habitat analyzed with quadrat method and composition of the plant in the deer diet using microhistological analysis of fecal sample. Forage species selection were assessed using a ivlev's index of electivity. Chemical properties analyzed were crude protein with Semi-micro Kjeldahl technique; gross energy content (GE) with a bomb calorimeter technique; ADF fibers with Goering and Van Soest method; mineral content of calcium (Ca) and phosphorus (P) by using the technique of Atomic Absobs Spectrophotometer; and tannin content was analyzed by *folin denish* reaction. The physical properties of food plants tested were water regain capacity and water solubility. Relationship between selection with chemical and physical properties of forage were determined by multiple regression statistical test. The results showed that feeding selection of timor deer are influenced by the chemical (content of CP, GE, ADF) and physical properties (water regain capacity and water solubility) of forage plants species. Feeding selection index increases with an increase in protein content, calcium content, the value of water regain capacity and water solubility of forage species. The content of CP and tannins, water regain capacity and water solubility of forage species can be a predictor variables to predict the feeding selection of forage species by timor deer in the habitat, with the regression model are as follows: $Y = -2.31 + 0.14X_1 + 0.04X_2 + 1.99X_3 - 0.12X_4$ (Y= selection index; X_1 = content of CP; X_2 = water regain capacity; X_3 = water solubility; and X_4 = content of tannin; $R^2 = 0.706$; $P < 0.05$).

Keywords: Timor deer, feeding selection, chemical properties, physical properties.

Introduction

Feeding selection on wildlife in nature is a complex problem, because it involves a variety of factors, both factors related to the nature of each type of plant feed (*physical, chemical and biological*), the availability of forage, disturbance factors or predators presence, social behavior, learning behavior and also a result of the evolutionary process (Hanley,1997; Moser *et al.*, 2006). Some research regarding the selection of plant species on timor deer (*Cervus timorensis*) in the wild indicate that deer are generalists or plants diversity of

feed high category that takes a lot of plants species, both grasses, forbs, as well as woody plants (Ginantra *et al.*, 2014; Allison, 2011; DeGarine-Wichatitsky *et al.*, 2005; Pattiselanno and Arobaya, 2009). Even the deer are also said to be opportunistic herbivores, because in scarce forage situations can also eat leftovers in the trash.

Research results about feeding selection of the timor deer (*C. timorensis*) in nature is more focused on the

availability of forage species in the habitat (DeGariné-Wichatitsky *et al.* (2005), Pattiselanno and Arobaya, 2009). Studies towards factors associated with physical properties (water regain capacity, water solubility), chemical properties (protein, energy, minerals, the content of the cell wall, and allelochemical) and biological properties (digestibility) of each species of forage plants have not been many do.

This study focuses on the relationship feeding selection on timor deer (*C. timorensis*) with factor of physical and chemical properties of each species of forage plants available in the habitat. Two of these properties are important to understand the role of nutritional factors in the feeding selection of animals

in natural habitat. An understanding of the selection of food plants and nutritional factors can be a reference in the management of the habitat to support the conservation of timor deer in Bali Barat National Park (BBNP).

Materials and Methods

Study Area

Field research was conducted in four habitat units on BBNP, namely; Cekik unit, Prapat Agung, Segara Rupek, and Brumbun (Figure 1), from January to March 2013 (rainy season) and July to September 2013 (dry season). Located at coordinates 8°11'15.14" - 8°06'21.13" S and 114° 26'25.45"- 114° 29'58.09" E, altitude between 4.90 to 90.84 m above sea level.




Figure 1. Location of study area at Bali Barat National Park

The type of vegetation in the habitat unit is savanna (Cekik, Segara Rupek and Brumbun) and Prapat Agung unit is a monsoon forest. Rainfall during the rainy season average of 220.17 mm/month and in the dry season average of 28.33 mm/month (data from BMKG region III Bali, 2013). Temperatures ranging from 24 °C to 37 °C and humidity ranging from 30 to 80%.

Analysis of Forage Species Composition in Habitat and in Deer Diet

Composition of plants species available in the habitat is determined by the quadrat method. Each quadrat

0.5 m x 0.5 m for grass/forbs vegetation, 1 m x 1 m for shrubs, and 5 m x 5 m for tree. Ten quadrats in each habitat unit applied per month, up to in each habitat unit used thirty quadrats per season. Percent ground cover each plant species measured in the quadrat. For shrubs and tree vegetation, percent cover of shoot that's count up to 1.2 m height (height level can be access by timor deer). The composition of each plant in the habitat (*ai*) is determined by the equation:

$$\frac{\text{average coverspecies-}i}{\text{total coverallsp}} \times 100\%$$

(average cover of species-*i* is the number of species-*i* cover divided by plot sampling) (Morrison, 2008).

Composition of plants species in the timor deer diet determined by techniques microhistological faecal sample (Holechek *et al.*, 1990). Fecal samples collected from five groups pellets at each habitat unit at 2-week intervals every month. Immediately collected fecal sample was dried (oven 70 °C) to avoid further decomposition. Samples on each habitat unit then compositing based on season to mikrohistology analysis. Identification of plant species in deer diets based on recognition of plant epidermic microscopic fragments preserved in the feces. Microscopic observations performed in the laboratory of plant taxonomy Udayana University of Bali. The composition of each plant in deer diets (ri) calculate by the formula:

$$\text{Composition of species-i} = \frac{\text{density species-i}}{\text{Total density all species}} \times 100\%$$

(density species-i is calculated based on the number of fragments of the epidermis of each species divided by the number slide total).

Forage selection were assessed using a ivlev's index of electivity (SI) (Krebs, 1989): $SI = (ri - ai) / (ri + ai)$, which is ai = composition (%) plant species-i in habitat, ri = composition (%) plant species-i in deer diet. SI values ranging from -1 to +1, where SI values 0.1 to 1 are indicate preference, 0.09 to -0.09 are proporsional, and -0.1 to -1 are avoidance.

Analysis of the Chemical and Physical Properties of Forage Plants

Samples of each species feed taken 250 grams of fresh ingredients in a period three times on each unit habitat. Each sample oven at 70 °C to obtain dry weight (DW). DW sample finely ground for analysis of chemical and physical properties. Crude protein (CP) analyzed by Semi-micro Kjeldahl technique, gross energy content (GE) by bomb calorimeter apparatus, fiber ADF (acid detergent fiber) analyzed by Goering and Van Soest method (Ranjhan and Krishna, 1980). Minerals (Ca and P) content were analyzed by atomic absorption spectrophotometer (Sinaga, 1997). Tannin content was analyzed by *folin denish* reaction (Suhardi, 1997).

The physical properties of each species of forage plants observed were water absorption (water regain capacity) and water solubility. Analysis procedure refers to Suhartati *et al.* (2004).

Statistical analysis

The relationship between the selection index with chemical properties (CP, Ca and P minerals, fiber ADF, tannins) and physical properties (water regain capacity and water solubility) of forage species used multiple regression statistical test. Test using the software "IBM SPSS 20".

Results and Discussion

Forage Selection by Timor Deer

There are 36 species were selected by the timor deer during the rainy season, which consists of 10 species of forbs, 18 species of graminoids, 8 species of woody plant. Thirty-eight species were selected in the dry season, which consists of 6 species of forbs, 17 species of graminoids, 15 species of woody plant. The same plant species were selected in the two seasons was found 26 species, which consists of 4 species of forbs, 16 species of graminoids, 6 species of woody plants. Nine species are selected only during the rainy season and 11 species were selected only in the dry season (Table 1).

Forage plant species on forbs category shows a high selection index in the two season. Based on selection index, nine of the ten species of this plant are selected in the rainy season preferred category (SI values 0.1 to 1). In the dry season, five of the six species of this plant are selected in preferred category. These species include *Commelina benghalensis*, *Boerhavia diffusa*, *Desmodium triflorum*, *Synedrella nodiflora*, and *Tribulus terrestris*. There are two species of plants are selected only in the dry season, namely *Vernonia patula* and *Justicia* sp.

In the category of graminoids, of eighteen species were selected in two seasons, 8 species is an important species for timor deer fodder in BBNP, because these species are preferred category (SI values 0.1 to 1). These species are; *Eriochloa ramosa*, *Axonopus compressus*, *Eriochloa subglabra*, *Dactyloctenium aegyptium*, *Panicum tryperon*, *Oplismenus Burmani*, *Digitaria adscendens*, *Eleusine indica*.

Some woody plant species the selection index is the fairly consistent in both seasons, including preference category as *Leucaena leucocephala*, *Grewia*

koordersiana and *Hibiscus tiliaceus*. In general, the selection of category woody plant selection index showed improvement in the dry season. There are five species of plants that are not selected in the rainy season, only are selected at the dry season with preferred categories, namely *Bridelia monoica*, *Gliricedia sepium*, *Zizyphus mauritiana*, *Pluchea indica* and *Sida acuta*. The species of forage on woody plant have become important as timor deer fodder when forbs and graminoids limited availability in the habitat.

These results indicate that the timor deer was able to accommodate the availability of the plants in quantity and quality. Certain plant species could become not preferred because of the presence of other species of preferred or a species to be favored than before because there are limited species in habitat. Selection behavior can change depending on the availability of habitat unit and season. Ginantra *et al.* (2014) and DeGariné-Wichatitsky *et al.* (2005) found that variations in the availability of plants from category forbs, graminoids and woody at the different types of habitat and season determine the feeding selection of timor deer. Hanley (1996) and Allison (2011), also stated that the availability of food plants in the habitat and season plays an important role in feeding selection behavior on deer.

Chemical Properties of Forage Species

Test of chemical properties (content of CP, GE, Ca, P, ADF and Total tannin), showing that the quality of the forage plants are available in BBNP is variaty. CP, mineral Ca and P content of forbs plants group is higher, but availability is high only during the rainy season. CP content of graminoids was lower, but GE content of its higher than forbs plants. CP content of woody plants and GE is high and availability is likely to be relatively longer than the forbs (Table 1). The same was reported by Amiri and Shariff (2012), Holechek (1984), that the content of the CP, GE and minerals vary based group of plants (forbs. Grasses, and woody plants).

Forage quality (in terms of the content of CP and GE) associated with the growth phase of a plant and environmental factors such as precipitation or soil moisture. The role of water is important for the metabolic process for the formation of organic compounds, so the content of CP and GE forage plants

in the rainy season becomes higher than the dry season. This statement is supported by some researchers that there is a seasonal dynamics of the nutrient content of forage in their natural habitat (rangelands) associated with the availability of water for plant growth (Rollins, 2011; Memmott *et al.*, 2011; Holechek, 1984). Further stated that the grasses and herbaceous dicots containing high nutrient levels when growth phase.

In general the tannin content of forage plants is still at a moderate level, which is less than 4% (Cannas, 2008). Tannin content of woody plants is relatively higher than forbs and graminoids. Holechek (1984), also reported that woody plants tend to contain a secondary metabolites including tannins higher than dicotyledonous herbaceous plants (forbs) and grasses category.

ADF content of forage species in forbs category is relatively lower than graminoids and woody plant category (Table 1). Results of this research is the logical consequence of lower ADF on forbs plants compared to others as a result of low fiber and high rough materials without nitrogen extract. Other researchers have also found that almost the same thing, namely grass (graminoids) is a group of plants that contain cell wall (ADF and NDF) tends to be higher than the herbaceous dicots (Amiri and Shariff, 2012; Holechek, 1984).

Physical Properties of Forage Species

Forbs and graminoids tend to decrease water absorption (water regain capacity) and water solubility in the dry season (Table 1). A decrease in physical properties is closely related to the aging phase of vegetative plant parts are accompanied by increased cell wall components (ADF and NDF). Water absorption and water solubility is influenced by the content of plant cell walls (fibers ADF), this statement can be proved by reference to the existence of a negative correlation between the content of ADF forage with water regain capacity and water solubility ($r = -0.450$ and $r = -0.632$) (Appendix 1 a). That is, the higher the content of the cell wall (ADF) plant, the water regain capacity and the solubility lower.

Unlike the case in woody plants, water absorption and water solubility (physical properties) relative almost the same in both seasons. Physiological conditions is due to the woody plants (shrubs and trees) plant parts

Table 1. Forage selection, chemical composition and physical properties of forage plants species in two season on BBNP (SI = selection index, CP= crude protein, GE= gross energy, Ca= Calcium, P= Phosphorus, ADF= acid detergent fiber, WRC= water regain capacity, and WS= water solubility)

No	Forage species	Rainy Season								Dry season							
		SI	CP %	GE MJ/kg	Ca %	P %	Tani n %	ADF %	WRC	WS	SI	CP %	GE MJ/kg	Ca %	P %	WRC	WS
	Forbs																
	<i>Commelina</i>																
1	<i>benghalensis</i>	0.63	22.44	14.74	2.29	0.46	0.33	28.25	7.89	0.41	0.58	13.54	13.53	2.01	0.40	4.71	0.42
2	<i>Boerhavia diffusa</i>	0.58	24.99	14.25	2.01	0.33	0.34	37.02	3.00	0.45	0.46	21.69	15.50	3.62	0.40	3.90	0.45
3	<i>Tribulus terrestris</i>	0.54	16.10	15.76	3.30	0.40	0.30	37.14	6.18	0.38							
4	<i>Desmodium triflorum</i>	0.51	21.92	16.48	1.48	0.45	0.52	35.17	6.29	0.47	0.24	17.23	17.66	1.52	0.16	3.84	0.42
5	<i>Fleura interupta</i>	0.50	16.36	17.52	1.50	0.38	0.61	42.77	5.63	0.43							
6	<i>Tephrosia pumila</i>	0.47	21.77	18.87	1.53	0.30	0.73	38.51	3.32	0.33	-	-	-	-	-	-	-
7	<i>Borreria laevis</i>	0.29	22.07	15.00	1.93	0.52	0.31	37.43	2.82	0.47	-	-	-	-	-	-	-
8	<i>Synedrella nodiflora</i>	0.28	21.25	14.68	1.95	0.71	0.39	46.34	2.38	0.48	0.56	20.13	14.89	2.56	0.51	3.41	0.38
9	<i>Acalypha indica</i>	0.28	19.81	14.80	3.87	0.59	0.47	34.05	3.75	0.47							
10	<i>Vernonia cinerea</i>	-0.24	16.13	13.97	2.09	0.14	0.43	44.71	3.22	0.51							
11	<i>Vernonia patula</i>										-0.06	14.24	18.67	1.58	0.35	3.19	0.26
12	<i>Justicia sp.</i>										0.50	17.05	13.65	4.75	0.25	4.10	0.33
	Graminoids/grasses																
1	<i>Eriochloa ramosa</i>	0.66	11.24	15.78	0.35	0.28	0.34	40.19	7.30	0.47	0.42	9.79	15.76	0.61	0.57	4.66	0.42
2	<i>Axonopus compresus</i>	0.65	15.46	17.44	0.44	0.27	0.30	44.97	6.71	0.36	0.38	12.77	16.91	0.53	0.25	5.73	0.33
2	<i>Eriochloa subglabra</i>	0.56	11.54	15.84	0.89	0.25	0.24	43.58	4.86	0.50	0.61	11.39	15.71	0.67	0.33	4.33	0.42
	<i>Dactyloctenium</i>																
4	<i>aegyptium</i>	0.53	13.44	15.63	1.13	0.27	0.35	44.19	5.54	0.35	0.31	10.6	15.62	0.75	0.29	3.57	0.43
5	<i>Panicum tryperon</i>	0.44	14.62	16.09	0.70	0.28	0.28	37.63	5.99	0.37	0.34	7.49	16.82	0.47	0.76	4.61	0.36
6	<i>Oplismenus burmani</i>	0.36	18.65	15.46	0.77	0.28	0.40	45.69	4.11	0.39	0.36	13.42	14.74	1.05	0.14	4.48	0.31
7	<i>Eleusine indica</i>	0.36	12.05	17.59	0.86	0.31	0.34	47.31	4.33	0.41	0.19	12.87	16.15	0.99	0.43	4.21	0.33
8	<i>Chloris barbata</i>	0.36	12.12	17.01	0.69	0.44	0.16	46.85	4.81	0.33	-0.04	9.6	16.12	0.59	0.40	3.72	0.27
9	<i>Cynodon dactylon</i>	0.34	14.94	18.09	0.59	0.32	0.37	41.16	4.84	0.37	0.05	8.99	17.14	0.64	0.65	4.23	0.36
10	<i>Digitaria adscendens</i>	0.21	11.19	17.50	0.56	0.27	0.32	48.30	4.72	0.32	0.38	12.07	16.27	0.82	0.35	3.98	0.36

Table 1. (continous)

No	Forage species	Rainy Season								Dry season							
		SI	CP %	GE MJ/kg	Ca %	P %	Tanin %	ADF %	WRC	WS	SI	CP %	GE MJ/kg	Ca %	P %	WRC	WS
11	<i>Panicum eruciforme</i>	-0.01	12.43	17.66	0.42	0.31	0.58	48.11	4.24	0.21	0.06	9.16	16.63	0.44	0.22	3.96	0.24
12	<i>Cyperus haspan</i>	-0.10	9.81	17.33	0.79	0.21	0.43	49.87	4.43	0.30							
13	<i>Imperata cylindrica</i>	-0.22	9.43	18.09	0.26	0.22	0.25	45.36	3.38	0.26	-0.29	7.20	17.03	0.32	0.16	3.57	0.28
14	<i>Themeda arguerns</i>	-0.23	7.40	17.33	0.21	0.20	0.34	52.54	3.78	0.23	-0.32	6.49	15.63	0.44	0.08	3.78	0.19
15	<i>Heteropogon contortus</i>	-0.28	9.04	17.96	0.46	0.23	0.61	49.52	3.39	0.27							
16	<i>Eragrostis amabilis</i>										0.14	9.57	17.39	0.56	0.24	4.71	0.39
17	<i>Andopogon aciculatus</i>	-0.29	9.98	17.04	0.75	0.30	0.41	48.02	3.18	0.21	-0.10	6.87	16.07	0.53	0.25	2.77	0.30
18	<i>Oplismenus composites</i>	-0.67	11.00	15.44	0.46	0.14	0.35	48.61	3.90	0.23	-0.03	12.67	15.69	0.24	0.15	4.26	0.27
19	<i>Phragmites sp.</i>	-0.85	8.47	18.70	0.36	0.26	0.29	51.58	2.72	0.23	-0.11	10.85	17.65	0.24	0.22	3.26	0.20
	Woody																
1	<i>Leucaena leucocephala</i>	0.51	27.65	20.22	1.21	0.27	2.16	27.62	4.74	0.46	0.51	21.28	19.28	1.91	0.23	4.19	0.47
2	<i>Hibiscus sinensis</i>	0.44	16.13	17.32	2.22	0.35	0.35	51.26	4.27	0.35	0.47	15.30	17.92	2.36	0.19	4.94	0.38
3	<i>Grewia koordersiana</i>	0.33	17.34	19.02	1.33	0.27	0.41	49.67	3.38	0.32	0.39	20.64	18.21	2.19	0.18	4.14	0.37
4	<i>Streblus asper</i>	0.17	15.22	15.94	1.82	0.11	0.35	50.45	2.62	0.26	-0.01	16.47	15.13	1.04	0.24	3.23	0.33
5	<i>Schleichera oleosa</i>	0.05	14.78	20.10	1.11	0.20	1.11	50.37	3.42	0.29	0.01	11.81	17.98	0.77	0.11	3.05	0.25
6	<i>Acacia auriculiformis</i>	0.00	21.14	22.05	0.69	0.10	0.51	64.23	2.39	0.23	0.13	22.79	21.16	0.75	0.11	2.71	0.31
7	<i>Malvastrum sp.</i>	-0.24	25.63	18.16	2.18	0.24	0.77	57.48	3.42	0.25	-	-	-	-	-	-	-
8	<i>Phylanthus emblica</i>	-0.60	11.97	19.73	0.60	0.07	2.75	41.01	2.42	0.36	-0.45	11.81	17.98	0.77	0.11	2.15	0.34
9	<i>Sida acuta</i>	-	-	-	-	-	-	-	-	-	0.53	19.08	16.52	2.43	0.56	3.98	0.33
10	<i>Bridelia monoica</i>	-	-	-	-	-	-	-	-	-	0.40	19.46	17.31	2.29	0.16	3.78	0.31
11	<i>Eupatorium odoratum</i>	-	-	-	-	-	-	-	-	-	-0.33	24.63	18.66	2.24	0.14	3.19	0.21
12	<i>Lantana camara</i>	-	-	-	-	-	-	-	-	-	-0.19	16.76	15.81	1.43	0.22	3.07	0.25
13	<i>Pluchea indica</i>	-	-	-	-	-	-	-	-	-	0.48	21.79	19.94	2.18	0.20	3.62	0.45
14	<i>Gliricedia sepium</i>	-	-	-	-	-	-	-	-	-	0.32	15.95	14.89	2.47	0.12	3.86	0.55
15	<i>Zizyphus mauritiana</i>	-	-	-	-	-	-	-	-	-	0.13	15.95	14.89	2.47	0.12	3.09	0.34
16	<i>Ipomoea hispida</i>	-	-	-	-	-	-	-	-	-	0.48	22.33	18.00	1.33	0.58	3.58	0.37

Note: The sign minus (-) in column = plant species exist in the habitat, but not selected ; blank column = species does not exist in the habitat

analyzed the shoots /young leaves (the part that still can eaten by deer), both in the rainy and dry seasons. On the other hand, the life of a plant or plant part young / old closely related to the content of its ADF and NDF. In line with Tillman *et al.* (1991) that the plants age or older leaves ADF and NDF content of its increasing or vice versa (positive correlation). Furthermore. if it is associated with the physical properties. especially its ability to hold water (water holding capacity) opposite occurs, ie ADF and NDF plant is negatively correlated with its physical properties (Behgar *et al.*, 2009).

Feeding Selection in Relation to the Chemical and Physical Properties of Forage

Feeding selection is influenced by several factors such as chemical feed content (CP, GE, minerals Ca, P, fiber ADF, tannins) and physical properties of forage plants (water regain capacity and water solubility). Results of this study are described with reference to the correlation between chemical factors, physical properties with feeding selection index.

Partial correlation (Pearson correlation) feeding selection with acontent of CP, GE. minerals Ca, mineral P, ADF, water regain capacity, and water solubility of forage plant species showed a significant correlation ($P < 0.05$), while the tannins are less significant correlation ($P > 0.05$).

The crude protein (CP) content of forage species positively contribute to feeding selection ($r = 0.457$; $P < 0.05$) (Table 2), increased CP content has positive effect on the level of selection. These results are in accordance with the feeding selection on sheep in the savannas that the content of CP on the species of forage on woody plants positively correlated with the selection of feed (Basha, 2012). Tixier *et al.* (2008) also found that feeding selection on red deer (*Cervus elaphus*) is positively correlated with protein content of feed on woody plants category in the spring, summer and autumn, and weak correlation to the category of grass plants. Selection relationship with a high protein content fodder grass category occurs in winter.

Water absorption (water regain capacity) contribute positively to the selection of forage ($r = 0.56$; $P < 0.05$). The species of plants that high water regain capacity show high level of selection (including the category of preferred), such as *Commelina*

benghalensis, *Tribulus terrestris* and *Desmodium triflorum* (forbs); *Eriochloa ramosa*, *Axonopus compresus*, *Eriochloa subglabra*, *Dactyloctenium aegyptium* (graminoids); *Leucaena leucocephala* and *Hibiscus tiliaceus* (woody plants). The ability to water absorb feed is one of the physical properties of the feed into the quality indicator of forage species, because it deals with digestibility, the higher the water absorption, the better the feed quality also. Results of this study are supported by Suhartati *et al.* (2004) that there is a positive relationship between the water absorption with the digestibility of feed. Behgar *et al.* (2009) also reported that the water absorption (water holding capacity) negatively correlated with the content of plant cell walls (ADF and NDF). This means, the lower the negative value of nutritional (such as ADF, NDF), the higher the water absorption. This relationship also explains that the level of plant feed digestibility (fermentative) higher as well.

Water solubility is positively correlated with the feeding selection ($r = 0.556$; $P < 0.05$) (Table 2). Water solubility of food plants also determine the level of digestibility of the feed. The higher the water solubility species of feed, feed digestibility simplify the process of mechanically through the role of saliva in the mouth. Thus. a very large contribution to the subsequent digestive process, both fermentative digestion in the rumen or in the post rumen hydrolytic. Kismunarto (2007) and Suhartati *et al.* (2004), stated that water solubility also determines the flow rate of feed to the post rumen, the higher the water solubility, the feed passes through the rumen faster.

The content of GE forage plant species showed a negative correlation with the selection of fodder ($r = -0.329$; $P < 0.05$) (Table 2). Physiologically, the selection can be approached from the deer eat eating voluntary aspect (VFI: Voluntary Feed Intake), which according to the Putra (1992) that VFI is affected by the GE content of the feed with a negative correlation. The content of DM was positively correlated with the content of GE feed. The higher DM content of GE feed means also higher. DM content and higher GE quicker to give a sense of satiety in the animal or animals faster fulfillment of energy and nutrients in general.

Minerals calcium (Ca) and phosphorus (P) contributed positively to the feeding selection ($r = 0.336$ and $r = 0.466$; $P < 0.05$) (Table 2). Relation of Ca and P minerals to feeding selection in the wild have not been

studied previously. The mineral content is Generally associated with the nutritional quality of forage species. Ceacero *et al.* (2010a) stated that, the deer could selecting mineral content in it's feed and the selection is formed by physiological effort to meet the need for minerals.

ADF contributes negatively to the feeding selection ($r = -0.503$; $P < 0.05$) (Table 2), increased content of ADF on the species of feed lowers feeding selection. ADF content associated with the feed palatability, the higher ADF means the palatability lower. The higher of ADF content can lower digestibility of feed, mean lower nutritive value of feed. Basha, (2012) and Hanley (1996) stated that the nutritive value of a species of feed determines the feeding selection in the wild herbivores, including deer.

Tannins showed a negative correlation with feeding selection, but weak correlation/non significant ($r = -0.235$; $P > 0.05$) (Appendix 1 a). Littlefield *et al.* (2011) and Lopez-Try *et al.* (2007) also found no significant correlation between tannin content with the feeding selection on white-tailed deer. The results showed that the tannin content of plant species on BBNP are at moderate levels (less than 4%), ranging from 0.16 to 2.75%. Cannas (2008) states that the tannins in moderate levels (less than 4%) do not cause negative effects for ruminants. Wheeler & Mochrie (1981) high tannin content can decrease the palatability of a species of feed that can affect the preferences of the type of feed. Some woody plants have tannin that can bind to proteins, thereby reducing the nutritive value of the feed.

Table 2. Correlations

		selectio	CP	GE	Ca	P	ADF	Tannin	WRC	WS
Pearson Correlation	selection	1.000	.457	-.329	.336	.466	-.503	-.235	.560	.556
	CP	.457	1.000	-.143	.592	.371	-.386	.245	-.058	.489
	GE	-.329	-.143	1.000	-.480	-.448	.433	.499	-.283	-.555
	Ca	.336	.592	-.480	1.000	.514	-.414	-.035	-.027	.520
	P	.466	.371	-.448	.514	1.000	-.474	-.306	.182	.523
	ADF	-.503	-.386	.433	-.414	-.474	1.000	-.159	-.450	-.632
	Tannin	-.235	.245	.499	-.035	-.306	-.159	1.000	-.239	.034
	WRC	.560	-.058	-.283	-.027	.182	-.450	-.239	1.000	.177
	WS	.556	.489	-.555	.520	.523	-.632	.034	.177	1.000
	Sig. (1- tailed)	selection	.	.003	.025	.023	.002	.001	.084	.000
CP		.003	.	.203	.000	.013	.010	.075	.369	.001
GE		.025	.203	.	.002	.003	.004	.001	.047	.000
Ca		.023	.000	.002	.	.001	.006	.419	.438	.001
P		.002	.013	.003	.001	.	.002	.035	.144	.001
ADF		.001	.010	.004	.006	.002	.	.177	.003	.000
Tannin		.084	.075	.001	.419	.035	.177	.	.081	.423
WRC		.000	.369	.047	.438	.144	.003	.081	.	.150
WS		.000	.001	.000	.001	.001	.000	.423	.150	.
N		selection	36	36	36	36	36	36	36	36
	CP	36	36	36	36	36	36	36	36	36
	GE	36	36	36	36	36	36	36	36	36
	Ca	36	36	36	36	36	36	36	36	36
	P	36	36	36	36	36	36	36	36	36
	ADF	36	36	36	36	36	36	36	36	36
	Tannin	36	36	36	36	36	36	36	36	36
	WRC	36	36	36	36	36	36	36	36	36
	WS	36	36	36	36	36	36	36	36	36

In this study, some plants such as *Phyllanthus Emblica*, *Leucaena leucocephala* and *Schleicera oleosa* which is species of plant that is eaten by the timor deer in BBNP, contain tannins which are relatively higher than other species, namely 2.75%, 2.16% and 1.11% respectively. Holechek (1984) states that the deer is one of the herbivores that have a smaller mouth parts which can be selectively utilize these plants with highly efficient, so that deer can minimize the concentration of tannins that enter the body. Nolte *et al.* (2004) stated that deer have the ability to eat plants that contain tannins, due to the mouth produces saliva proteins capable of binding tannins and minimize the damaging effects of tannins on the digestibility.

Results of regression correlation test (stepwise method) between the feeding selection with the

content of the CP, GE, Ca, P, ADF, tannins, water absorption and water solubility of forage species, it feeding selection showed a significant regression correlation with four predictor variables. Four variables are water regain capacity (X_1), the content of CP (X_2), water solubility (X_3) and the content of tannins (X_4). Regression model of feeding Selection with four predictor variables are as follows: $Y = -2.31 + 0.14X_1 + 0.04X_2 + 1.99X_3 - 0.12X_4$ (Table 3). With R determinant coefficient value (R^2) is 0.706, meaning that jointly four factors (water regain capacity, content of CP, water solubility and tannin content influential of 70.6% against the selection forage plant species in the habitat, and the remainder was due to other factors.

Table 3. Model Summary

Model	R	R Square	Adjusted R	Std. Error of the	Change Statistics				
					R Square	F Change	df1	df2	Sig. F
1	.560 ^a	.313	.293	.33624	.313	15.502	1	34	.000
2	.744 ^b	.553	.526	.27518	.240	17.762	1	33	.000
3	.785 ^c	.616	.580	.25917	.062	5.202	1	32	.029
4	.814 ^d	.662	.619	.24685	.047	4.274	1	31	.047
5	.840 ^e	.706	.657	.23423	.043	4.432	1	30	.044

1. Predictors: (Constant), WRC; 2. Predictors: (Constant), WRC, CP; 3. Predictors: (Constant), WRC, CP, WS; 4. Predictors: (Constant), WRC, CP, WS, Tannin; 5. Predictors: (Constant), WRC, CP, WS, Tannin, GE

Coefficients^a

Model		Unstandardized Coefficients		Standardized Coefficients	t	Sig.	Correlations			Collinearity Statistics	
		B	Std. Error				Beta	Zero-order	Partial	Part Tolerance	VIF
		1	(Constant)	-.506	.182		-2.782	.009			
	WRC	.158	.040	.560	3.937	.000	.560	.560	.560	1.000	1.000
2	(Constant)	-	.212		-5.386	.000					
	WRC	.166	.033	.588	5.046	.000	.560	.660	.587	.997	1.003
	CP	.044	.010	.491	4.215	.000	.457	.592	.490	.997	1.003
3	(Constant)	-1.327	.216		-6.157	.000					
	WRC	.149	.032	.527	4.668	.000	.560	.636	.511	.941	1.063
	CP	.031	.011	.343	2.693	.011	.457	.430	.295	.739	1.353
	WS	1.266	.555	.295	2.281	.029	.556	.374	.250	.718	1.393
4	(Constant)	-1.069	.240		-4.452	.000					
	WRC	.135	.031	.478	4.342	.000	.560	.615	.453	.898	1.114
	CP	.036	.011	.403	3.228	.003	.457	.502	.337	.700	1.429
	WS	1.212	.529	.282	2.288	.029	.556	.380	.239	.716	1.396
	Tannin	-.068	.033	-.229	-2.067	.047	-.235	-.348	-	.887	1.127

	(Constant)	- 2.311	.632		-3.655	.001					
	WRC	.141	.030	.499	4.751	.000	.560	.655	.470	.890	1.124
5	CP	.036	.011	.400	3.383	.002	.457	.526	.335	.700	1.430
	WS	1.990	.624	.463	3.190	.003	.556	.503	.316	.465	2.152
	Tannin	-.116	.039	-.390	-3.000	.005	-.235	-.480	-	.580	1.724
	GE	.000	.000	.321	2.105	.044	-.329	.359	.208	.420	2.378

a. Dependent variable: selection index

Selection of forage plants on deer in the wild is a complex behavior and a consequence of interactions among several factors. Selection of the species of feed on deer involves several factors, such as nutritional value, alelokimia compound (negative value), the physical properties of the plant itself, the availability in habitat and also for competitors or predators. Likewise, olfactory and taste organs of animals may be used to determine chemical compound or nutrient which is essential for life. Given the social learning process, learning from the mother and the process of evolution has formed a feeding selection behavior in animals in nature (Ceacero *et al.*, 2010b; Hanley, 1997; Moser *et al.*, 2006).

Conclusion

Feeding selection of forage plants in the timor deer are influenced by the chemical (content of CP, GE, ADF) and physical properties (water regain capacity and water solubility) species of forage plants. Feeding selection increases with an increase in protein content, calcium content, the value of water regain capacity and water solubility of forage species. The content of CP and tannins (chemical properties), water regain capacity and water solubility (physical properties) of forage plants species can be a predictor variables to predict the feeding selection of forage plants in the habitat, with the regression model are as follows: $Y = -2.31 + 0.14X_1 + 0.04X_2 + 1.99X_3 - 0.12X_4$ (X_1 = content of CP; X_2 = water regain capacity; X_3 = water solubility; and X_4 = content of tannin; $R^2 = 0.706$; $P < 0.05$).

Acknowledgments

Thanks to the Research Center and Community Services of Udayana University who has funded this study. Thank you to the hall manager, staff of BBNP over all facilities for the conduct of research and helped the collection of samples in the field.

References

- Allison, K. 2011. Feeding Behavior of Deer. Available from : <http://www.ehow.com/search.html>.
- Amiri, F and Shariff, A.R.B.S.M. 2012. Comparison of nutritive values of grasses and legume species using forage quality index. *Songklanakarin J. Sci. Technol.* 34 (5), 577-586.
- Basha, N.A.D. 2012. Feeding behaviour, diet selection of goats and nutritive value of browse species in sub-humid subtropical savannah]. (Dissertation). School of Agricultural, Earth and Environmental Sciences, University of KwaZulu-Natal, Pietermaritzburg, South Africa.
- Behgar, M., Valizadeh, R., Mirzaee, M., Naserian, A.A. and Nasiri, M.R. 2009. Correlation Between the Physical and Chemical Properties of Some Forages and Non-Forage Fiber Sources. *Journal of Animal and Veterinary Advances*, Volume: 8 Issue: 11 Page No.: 2280-2285.
- Ceacero, F., Castillejos, T.L., García, A.J., Estévez, J.A. and Laureano, G. 2010a. Physiological variables explain mineral intake in Iberian red deer. *Physiology & Behavior* 100 (2010) 122–127.
- Ceacero, F., Castillejos, T.L., García, A.J., Estévez, J.A. and Laureano, G. 2010b. Can Iberian red deer (*Cervus elaphus hispanicus*) discriminate among essential minerals in their diet? *British Journal of Nutrition* (2010), 103, 617–626.
- Cannas, A. 2008. Tannins: fascinating but sometimes dangerous molecules. Cornell University. Department of Animal Science. Available from: <http://www.ansci.cornell.edu/plants/toxicagents/tannin.html>
- DeGariné-Wichatitsky, M., Soubeyran, Y., Maillard, D. and Duncan, P. 2005. The Diets of introduced Rusa Deer (*Cervus timorensis russa*) in a Native Sclerophyll Forest and a Native Rainforest of New Caledonia. *New Zealand Journal of Zoology*, Vol. 32:117-126.

- Ginantra, K., S. Putra, W. Suarna and W. Kasa. 2014. Botanical Composition of forage by Timor Deer (*Cervus timorensis* Blainville) in A Monsoon Forest and Savanna of West Bali National Park. *Int. J. Pure App. Biosci.* 2 (5): 205-213 (2014).
- Hanley, T.A. 1997. A Nutritional View of Understanding and Complexity in the Problem of Diet Selection by Deer (Cervidae). *Oikos* 79:209-218, Copenhagen.
- Holechek, J.L., Vavra, M. and Pieper, R.D. 1990. Methods for Determining the Botanical Composition, Similarity, and Overlap of Range Herbivore Diets. National Research Council/National Academy of Sciences, London.
- Holechek, J.L. 1984. Comparative Contribution of Grasses, Forbs, and Shrubs to the Nutrition of Range Ungulates. *Rangelands* 6 (6).
- IBM SPSS. 2011. Statistical Product and Service Solutions (SPSS 20). IBM Corporation.
- Krebs, J.C. 1989. Ecological Methodology. Harper Collins Publishers, New York.
- Littlefield, K.A., Mueller, J.P., Muir, J.P. and Lambert, B.D. 2011. Correlation of Plant Condensed Tannin and Nitrogen Concentrations to White-Tailed Deer Browse Preferences in the Cross Timbers. *The Texas Journal of Agriculture and Natural Resource* 24:1-7.
- Memmott, K.L., Anderson, V.J. and Fugal, F. 2011. Seasonal Dynamics of Forage Shrubs Nutrients. *Rangelands*, 33 (6):12-16.
- Morrison, J.I., 2008. Using Microhistological Techniques to Predict Botanical Composition of Horse Diets on Cool-Season Grass pasture. (A thesis submitted) College of Agriculture at The University of Kentucky.
- Moser, B., Schütz, M. and Hindenlang, K.E. 2006. Importance of alternative food resources for browsing by roe deer on deciduous trees: The role of food availability and species quality. *Forest Ecology and Management* (226): 248–255. Available from: www.sciencedirect.com.
- Nolte, D.L., Kimbal, B.A., Perry, K.R., Villalba, J.J. and Provenza, F.D. 2004. Effects of Forage Nutritional Quality (Energy and Protein) on Deer Acceptance of Foods Containing Secondary Metabolites. *USDA National Wildlife Research Center*. Available from : http://digitalcommons.unl.edu/icwdm_usdanwrc/374.
- Pattiselanno, F. and Arobaya, A.Y.S. 2009. Grazing Habitat of the Rusa Deer (*Cervus timorensis*) in the Upland Kebar, Manokwari. *Jurnal Biodiversitas* ISSN: 1412-033X (printed edition), Volume 10, Number 3, July 2009 ISSN: 2085-4722 (electronic), Pages: 134-138.
- Putra, S. 1992. Evaluation of Chemical Composition and Feed Intake of 16 Provenan Gamal (*Gliricidia sepium*) that Planted on Dryland in Bali (Thesis). Bogor: Bogor Institute of Agriculture.
- Ranjhan, S.K and G. Krishna. 1980. *Laboratory Manual for Nutrition Research*. Vikas Publishing House Pvt. Ltd., New Delhi.
- Rollins, D. 2011. Nutritional Value of Deer Habitat. Texas Natural Wildlife. Available from: <http://agrillife.org/texnatwildlife/deer/factors-effecting-deer-diets-and-nutrition1-2393/nutritional-value-of-deer-habitat/>
- Sinaga, Y. 1997. Elephant grass as Substitute Control Macro Mineral Analysis on Forage. Livestock research centers Ciawi, Bogor.
- Suhartati, F.M., Suryapratama, W. dan Rahayu, R. 2004. Analysis of the Physical Properties of a Local Grass s. *Animal Production*, Vol. 6 No: 1: 37-42
- Suhardi, 1997. Polyphenol Compounds Analysis Products Fruits and Vegetables. Vol.3. Processing Biochemistry Laboratory, Faculty of Agricultural Technology. Gadjah Mada University, Yogyakarta.
- Tixier, H.V., Renaud, P.C., Morellet, N., Jamot, J., Besle, J.M. and Dumont, B. 2008. Selection for Nutrients by Red Deer Hinds Feeding On Mixed Forest Edge *Oecologia*, Springer-Verlag, 156: 715-726.
- Wheeler, J.L. and Mochrie, R.D. 1981. Forage Evaluation: Concepts and Techniques. American Forage and Grassland Council, CSiro.